

Blue lines indicate the area meeting the ISRA Criteria; dashed lines indicate the suggested buffer for use in the development of appropriate place-based conservation measures

PETRIE ISRA

New Zealand & Pacific Islands Region


SUMMARY

Petrie is located on the northeastern side of New Caledonia. The area is an atoll with a barrier reef surrounding an inner lagoon that opens to oceanic waters on the northwestern side. It is characterised by live and dead coral substrates with sandy patches. The area overlaps with the Natural Park of the Coral Sea. Within this area there are: **threatened species** and **undefined aggregations** (Grey Reef Shark Carcharhinus amblyrhynchos).

-	-
NEW CALEDON	IA
-	-
0-50 metre	es
-	-
48.55 km²	

CRITERIA

Criterion A - Vulnerability; Sub-criterion C5 - Undefined Aggregations



sharkrayareas.org



DESCRIPTION OF HABITAT

Petrie is located on the northeastern side of New Caledonia within the Coral Sea. It sits on the most northwestern part of the Loyalty Ridge, ~180 km of the northern tip of the Grande Terre Island and ~290 km northwest of Ouvéa Island (Maurizot et al. 2020). The area is an atoll with a barrier reef surrounding an inner lagoon that opens to oceanic waters on the northwestern side (Andréfouët & Torres-Pulizza 2004). It is characterised by a narrow barrier reef with a steep reef slope dominated by live and dead coral substrates along with sandy and rocky patches (Pelletier et al. 2020).

The area overlaps with the Natural Park of the Coral Sea (UNEP-WCMC & IUCN 2024).

This Important Shark and Ray Area is benthic and pelagic and is delineated from surface waters (O m) to 50 m based on the bathymetry of the area.

ISRA CRITERIA

CRITERION A - VULNERABILITY

One Qualifying Species considered threatened with extinction according to the IUCN Red List of Threatened Species regularly occur in the area. This is the Endangered Grey Reef Shark (Simpfendorfer et al. 2020).

SUB-CRITERION C5 - UNDEFINED AGGREGATIONS

Petrie is an important area for undefined aggregations of one shark species.

Aggregations of Grey Reef Sharks are regularly observed in this area and have been documented by underwater surveys, visual censuses, and Baited Remote Underwater Video Station (BRUVS) surveys (Vigliola et al. 2014; Schohn et al. 2017; A Friedlander unpubl. data. 2024; Wantiez et al. unpubl data 2024).

Between 2012-2014, underwater surveys (n = 253) were conducted along New Caledonia reefs. Surveys consisted of fixed-width transects (25 m long by 4 m wide at 10 and 20 m depths), distancesampling visual census (50 m long by 10 m wide at ~15 m depth), and BRUVS (1 hour at ~20 m depth; Vigliola et al. 2014). Of these surveys, 18 were conducted in Petrie during 2013, showing that Grey Reef Sharks aggregate in the area (Vigliola et al. 2014). Carcharhinids were observed in 89% of the dives with Grey Reef Shark recorded in 78% of the surveys, including aggregations (mean = 9, maximum = 20 individuals; Vigliola et al. 2014). In addition, surveys with BRUVS (2012-2014) and distance-sampling visual census (2012-2014) were conducted at 385 sites for BRUVS and 2,790 sits for visual census (including 15 marine protected area) along New Caledonia with Petrie holding one of the largest abundances for sharks among all sites (Juhel et al. 2017). Aggregations for Grey Reef Sharks in visual censuses (8-14 individuals) were among the largest in all New Caledonia (Juhel et al. 2017). Further, of 34 surveys with rotating video stations conducted in 2014 in the area (deployed for 12 minutes from the boat and consisting of cameras rotating 60° every 30 seconds), Grey Reef Sharks were observed in ~40% of the deployments (Schohn et al. 2017). More information is needed to confirm the nature and function of these aggregations.

Acknowledgments

Alan Friedlander (National Geographic/Pristine Seas Project), Laurent Wantiez (Université de la Nouvelle-Caledonie), Laurent Vigliola (Institut de recherche pour le développement), Jean-Baptiste Juhel (Centre national de la recherche scientifique, Université de Guyane), and Emiliano García-Rodríguez (IUCN SSC Shark Specialist Group – ISRA Project) contributed and consolidated information included in this factsheet. We thank all participants of the 2024 ISRA Region 10 – New Zealand and Pacific Islands workshop for their contributions to this process.

This factsheet has undergone review by the ISRA Independent Review Panel prior to its publication.

This project was funded by the Shark Conservation Fund, a philanthropic collaborative pooling expertise and resources to meet the threats facing the world's sharks and rays. The Shark Conservation Fund is a project of Rockefeller Philanthropy Advisors.

Suggested citation

IUCN SSC Shark Specialist Group. 2024. Petrie ISRA Factsheet. Dubai: IUCN SSC Shark Specialist Group.

QUALIFYING SPECIES

Scientific Name	Common Name	IUCN Red List Category	Global Depth Range (m)	ISRA Criteria/Sub-criteria Met								
			•	Α	В	Cı	C2	C3	C4	C5	Dı	D2
SHARKS												
Carcharhinus amblyrhynchos	Grey Reef Shark	EN	0-280 m	Х						Х		



SUPPORTING SPECIES

Scientific Name	Common Name	IUCN Red List Category
SHARKS		
Triaenodon obesus	Whitetip Reef Shark	VU

IUCN Red List of Threatened Species Categories are available by searching species names at <u>www.iucnredlist.org</u> Abbreviations refer to: CR, Critically Endangered; EN, Endangered; VU, Vulnerable; NT, Near Threatened; LC, Least Concern; DD, Data Deficient.





REFERENCES

Andréfouët S, Torres-Pullizza D. 2004. Atlas des récifs coralliens de Nouvelle-Calédonie. Nouméa: Institut de Recherche pour le développement-IFRECOR.

Juhel JB, Vigliola L, Mouillot D, Kulbicki M, Letessier TB, Meeuwig JJ, Wantiez L. 2017. Reef accessibility impairs the protection of sharks. *Journal of Applied Ecology* 55: 673–683. https://doi.org/10.1111/1365-2664.13007

Maurizot P, Collot J, Cluzel D, Patriat M. 2020. Chapter 6 The Loyalty Islands and Ridge, New Caledonia. In: Maurizot P, Mortimer N, eds. *New Caledonia*: Geology, Geodynamic Evolution and *Mineral Resources*. London: The Geological Society of London, 131–145 pp. https://doi.org/10.1144/M51-2017-24

Schohn T, Pelletier D, Carpentier L. 2017. Etat de santé des habitats et peuplements de poissons des îles et récifs éloignés Astrolabe, Pétrie et Walpole Parc Naturel de la Mer de Corail - Evaluation initiale par stations vidéo rotatives STAVIRO 2014. Nouméa: IFREMER.

Simpfendorfer C, Fahmi, Bin Ali A, Utzurrum JAT, Seyha L, Maung A, Bineesh KK, Yuneni RR, Sianipar A, Haque AB, et al. 2020a. Carcharhinus amblyrhynchos. The IUCN Red List of Threatened Species 2020: e.T39365A173433550. https://dx.doi.org/10.2305/IUCN.UK.20203.RLTS.T39365A173433550.en

UNEP-WCMC & IUCN. 2024. Protected Planet: The World Database on Protected Areas (WDPA) and World Database on Other Effective Area-based Conservation Measures (WD-OECM) [Online], February 2024, Cambridge, UK: UNEP-WCMC and IUCN. Available at: www.protectedplanet.net Accessed Octobre 2024.

Vigliola L, Wantiez L, Kulbicki M, Ballesteros E, Brown E, D'Agatha S, Friedlander A, Gossuin H, Juhel JB, Mouillot D, Sala E. 2014. Pristine Mer de Corail: Les récifs éloignés de Nouvelle-Calédonie. Rapport au Gouvernement et aux trois Provinces de la Nouvelle Calédonie. Nouméa: Institut de Recherche pour le développement-Université de la Nouvelle-Calédonie-Université de Montpellier 2-National Geographic Society.