

Blue lines indicate the area meeting the ISRA Criteria; dashed lines indicate the suggested buffer for use in the development of appropriate place-based conservation measures

PIURA-LA LIBERTAD ISRA

Central and South American Pacific Region

SUMMARY


Piura-La Libertad extends from the southern Piura region to northern La Libertad region in Peru. It is located within the Northern Humboldt Current System, one of the most productive marine systems in the world and overlaps with major upwelling centres. The area has exceptional marine productivity and a variety of habitats dominated by rocky shores and sandy substrates. It overlaps with Lobos de Tierra and Lobos de Afuera islands, a marine protected area system which are both Key Biodiversity Areas. Within this area there are: **threatened species** (e.g., Spotted Houndshark *Triakis maculata*); **range-restricted species** (e.g., Humpback Smoothhound *Mustelus whitneyi*); **reproductive areas** (e.g., Smooth Hammerhead *Sphyrna zygaena*); **feeding areas** (e.g., Chilean Eagle Ray *Myliobatis chilensis*); and the area sustains a **high diversity of species** (17 species).

CRITERIA

Criterion A - Vulnerability; Criterion B - Range Restricted;
Sub-criterion C1 - Reproductive Areas; Sub-criterion C2 - Feeding Areas
Sub-criterion D2 - Diversity

—	—
PERU	—
—	—
0-1,000 metres	—
—	—
39,187 km²	—
—	—





DESCRIPTION OF HABITAT

Piura-La Libertad extends from the southern Piura region to northern La Libertad region in Peru. The area overlaps with the Northern Humboldt Current System (NHCS), which is one of the most productive ocean ecosystems due to its coastal upwelling producing a high abundance of zooplankton that sustains the ecosystem (Pennington et al. 2006). Oceanographic features, associated with wind forcing, create strong upwelling and high levels of primary productivity in the offshore waters of northern Peru (Bakun & Weeks 2008; Montecino & Lange 2009). Warm subtropical waters move closer to the coast in the austral summer and autumn, and coastal upwelling disperses them in winter and spring (Bakun & Weeks 2008). This upwelling, and its associated productivity, allow the development of an exceptionally high biomass of ecologically important marine species such as Peruvian Anchoveta *Engraulis ringens* and Humboldt Squid *Dosidicus gigas* (Bakun & Weeks 2008; Gonzalez-Pestana et al. 2022).

Within the NHCS, six major upwelling centres have been identified which are characterised by a higher concentration of phytoplankton. These centres tend to be associated with prominent continental features such as peninsulas and semi-protected bays where nutrients are concentrated, producing ‘upwelling shadows’ (Graham et al. 1992).

This area overlaps with Punta Illescas which represents the northern most upwelling centre. Punta Illescas has unique coastal topography as the second westernmost point of the south-eastern Pacific shoreline and produces a strong upwelling (Chavez & Messie 2009). It is also at the narrowest part of the continental shelf along the Peruvian coast which enhances productivity (Jacox & Edwards 2011). The northern area is influenced by the Cromwell Current which is the main contributor to coastal upwelling in northern Peru (Zuta & Guillen 1970), allowing a rich demersal subsystem due to its high oxygen levels (Vargas & Mendo 2010), supporting the Peruvian Hake *Merluccius gayi peruanus* demersal fishery.

This area overlaps with Lobos de Tierra and Lobos de Afuera islands which are part of Sistema de Islas, Islotes y Puntas Guaneras marine protected area system (UNEP-WCMC & IUCN 2026), and have been identified as Key Biodiversity Areas (KBA 2026). This area also overlaps with two Ecologically or Biologically Significant Marine Areas (EBSA): the Northern Humboldt Current System EBSA and the Major Upwelling Centers and Seabird Associated with the NHCS EBSA (CBD 2026).

This Important Shark and Ray Area is delineated from surface waters (0 m) to 1,000 m based on depth of the Qualifying Species in the area.

ISRA CRITERIA

CRITERION A – VULNERABILITY

Sixteen Qualifying Species considered threatened with extinction according to the IUCN Red List of Threatened Species regularly occur in the area. Threatened sharks comprise four Critically Endangered, two Endangered, and four Vulnerable species. Threatened rays comprise five Vulnerable species, and one Critically Endangered (IUCN 2025).

CRITERION B – RANGE RESTRICTED

Piura-La Libertad holds the regular presence of six resident range-restricted species: Chilean Angelshark, Humpback Smoothhound, Spotted Houndshark, Chilean Eagle Ray, Peruvian Eagle Ray, and Shorttail Fankate. Chilean Angelshark, Spotted Houndshark, Chilean Eagle Ray, and Peruvian

Eagle Ray are restricted to the Humboldt Current LME. Shorttail Finskate occurs primarily in the Humboldt Current LME and only marginally into the Pacific Central-American Coastal LME. Humpback Smoothhound occurs in the Humboldt Current LME and the Pacific Central-American Coastal LME.

These species are regularly encountered and often targeted in small-scale fisheries (Alfaro-Shigueto et al. 2010; Céspedes 2014; Gonzalez-Pestana et al. 2016, 2022; Córdova-Zavaleta 2022; Sencio-Sánchez et al. 2025; Garnique-Capuñay et al. 2025). Some of them represent the most landed sharks (Chilean Angelshark, Humpback Smoothhound) or rays (Chilean Eagle Ray, Peruvian Eagle Ray) in Peru and have their most important landing sites and fishery areas in this area (Gonzalez-Pestana et al. 2016, 2022; IMARPE landings statistics between 2010-2020). Shorttail Finskate is the most abundant (according to biomass) bycatch ray species in the Peruvian Hake industrial trawling fishery which operates within this area (Céspedes 2014).

SUB-CRITERION C₁ – REPRODUCTIVE AREAS

Piura-La Libertad is an important reproductive area for five shark species.

The Peruvian fishery for Smooth Hammerhead that operates in this area is composed of neonates, young-of-the-year, juveniles, and adult females (Castañeda 2001; Gonzalez-Pestana 2014, 2018; Torres 2018; Córdova-Zavaleta 2022). Studies with samples from ~4,000 individuals indicate that the smallest individuals measured 44 cm of total length (TL) while the size ranged between 70-90 cm TL. Size-at-birth for this species is reported at 49-63 cm TL (Rigby et al. 2019). During late austral spring and early summer (November to January) females in advanced pregnancy stages were captured as this life-stage is targeted (Castañeda 2001; Gonzalez-Pestana 2014, 2018). Sharks are born in late spring and early summer (with open umbilical scars recorded), and they stay in this area for their first year with some individuals staying up to two years (Gonzalez-Pestana 2014, 2018).

Tope Shark is caught by small-scale fisheries operating in this area. San Jose and Santa Rosa (Lambayeque region) are the most important landing sites for this species along the Peruvian coast, representing 78% of the total landings of the species in Peru (IMARPE national reports). Between 2015-2019, 382 individuals were sampled within this area. In San Jose (Lambayeque region), animals measured on average 112.5 ± 29 cm TL with a minimum size of 50 cm TL, and in Salaverry (La Libertad region) an average 132.1 ± 23.2 cm TL with a minimum size of 80 cm TL (Córdova-Zavaleta 2022). Most individuals were juveniles (size-at-maturity: 206-235 [males] and 227-244 [females] cm TL; Ebert et al. 2021), and some were young-of-the-year since size-at-birth is between 26-40 cm TL.

Between 2015-2019, 954 Copper Sharks were sampled measuring 105.6 ± 22.3 cm TL with a minimum size of 50 cm TL (Córdova-Zavaleta 2022). Most individuals were juveniles (size-at-maturity: 120-135 [males] and 134-140 [females] cm TL; Peres & Vooren 1991), and some were neonates and young-of-the-year since size-at-birth is between 59-70 cm TL (Ebert et al. 2021; Drew et al. 2017).

Pregnant females of Humpback Smoothhound have been observed in this area. Sampled individuals (n = 41) from March to July 2013, and May to September 2016, were pregnant females (average body size: 88.1 ± 17.8 cm TL) in varied stages of embryonic development. Neonates (n = 16) were also recorded (Gonzalez-Pestana et al. 2019). The largest embryo measured 23 cm TL and the smallest free-living neonate, with an open umbilical scar, measured 22.4 cm TL. Additionally, captures of gravid females have been observed during two fishery trips off northern Lambayeque coast and off Punta Illescas (Adriana Gonzalez-Pestana pers. obs. 2022).

Eighty to 85% of Chilean Eagle Ray landings within this area were composed of immature individuals, including neonates (with minimum body sizes of 30 cm disc width [DW]), with mature adult females (some pregnant with litter sizes of 2–4 pups commonly captured in the austral summer; Torres 1978; Castañeda 1994). Between 2015–2019, 4,577 individuals were sampled measuring on average 75.6 ± 28.7 cm DW with minimum sizes of 20 cm DW (Córdova-Zavaleta 2022). For this species, the size-at-maturity is 115 cm DW (Castañeda 1994). The size-at-birth is unknown; yet embryos with a body size of 29 cm DW have been reported in this area (Castañeda 1994). Thus, mostly juveniles, including neonates, are recorded in this area.


SUB-CRITERION C2 – FEEDING AREAS

Piura-La Libertad is an important feeding area for eight sharks. The two most common prey species (Peruvian Anchoveta and Humboldt Squid) represent one of the most abundant marine resources worldwide as these are the most caught species (fish and invertebrate, respectively) (FAO 2022). Peru is responsible for the largest volumes captured for both species (FAO 2022). The Peruvian Anchoveta is one of the main reasons why the NHCS produces more fish per surface unit than any other marine ecosystem (Chavez et al. 2008). Within Peru, historically the largest fishery has been concentrated in northern Peru for Humboldt Squid (Csirke et al. 2018) and in northern-central Peru for Peruvian Anchoveta (Castillo et al. 2015), both located within this area. During warm periods (El Niño events or summers), these species aggregate in major upwelling centres associated with the NHCS EBSA (with traditional ecological evidence that Humboldt Squid aggregates in the central part of this area), as these centres serve as a refuge, given the persistence of upwelling in them (Bertrand et al. 2004, CBD 2017, Jian et al. 2020). Also, in the northern part of this area, one of the largest abundances of Peruvian Hake has been recorded as the fishery has developed into one of the main fisheries in Peru (Arellano & Swartzman 2010).

Results from a diet analysis of 485 Smooth Hammerheads (neonates, young-of-the-year, juveniles, and adult females) between 2013–2015, found that Humboldt Squid (27% Index of Relative Importance [IRI]) and Patagonian Squid *Doryteuthis gahi* (37% IRI) were the main prey (Gonzalez-Pestana et al. 2017). Smooth Hammerheads presented a narrow trophic niche (i.e., highly specialised predator) in this area. In this study, 78% of stomachs contained food items and in one adult (230 cm TL) stomach, 74 pairs of squid beaks were counted (the equivalent of 74 cephalopods) (Gonzalez-Pestana et al. 2017). Other diet studies of Smooth Hammerhead in the eastern Pacific Ocean (Ecuador and Baja California) indicate that this species feeds mainly on cephalopods (Bolaño 2009; Estupiñan-Montaño et al. 2009; Galvan-Magaña et al. 2013). Smooth Hammerhead has been the third most captured shark species by fisheries in Peru and the most frequently captured shark species off northern Peru between 1997–2021 (Gonzalez-Pestana et al. 2016; IMARPE landings statistics). Most of the landings and fishing areas are in Piura, Lambayeque, and La Libertad (within this area). One of the most important fishing areas in Peru are around the Lobos de Tierra and Lobos de Afuera Islands, offshore of Lambayeque (Carbajal et al. 2007; Llanos et al. 2009). Thus, this area represents an important feeding ground for this species along the Eastern Pacific.

An analysis of stomach contents from 44 Tope Sharks (adults and juveniles) between January 2015 and August 2016, determined that it preys mostly on teleost fishes (Peruvian Hake and Peruvian Anchoveta) and secondarily on cephalopods (Gonzalez-Pestana et al. 2021a).

An analysis of stomach contents from 69 Copper Sharks (adults and juveniles) between January 2015 and August 2016, determined that it preys mostly on teleosts in which the Peruvian Anchoveta is the most important prey (43% Prey-specific Index of Relative Importance [PSIRI]). This species is considered a predator with a high degree of specialisation (Gonzalez-Pestana et al. 2021a). In other



places where its diet has been studied (South Africa and Argentina), its diet is composed mainly of small pelagic schooling fishes (Cliff & Dudley 1992; Lucifora et al. 2009; Smale 1991).

An analysis of stomach contents from 76 Humpback Smoothhounds (mostly adults) between January 2015 and August 2016, determined that the species preys mainly on Peruvian Anchoveta (21% PSIRI), and secondarily on crustaceans (crabs and stomatopods) and molluscs (gastropods and cephalopods) (Samame et al. 1989; Gonzalez-Pestana et al. 2021a). Diet varies according to season (Samame et al. 1989).

An analysis of stomach contents from 43 Spotted Houndsharks between January 2015 and August 2016, determined that the species preys mostly on teleost fishes (e.g., Peruvian Anchoveta) and crustaceans (Gonzalez-Pestana et al. 2021a).

An analysis of stomach contents from 72 Broadnose Sevengill Sharks between 2015 and 2019, determined that the species preys mostly on teleost fishes and marine mammals (sea lions and small cetaceans). This diet varies according to ontogeny with larger individuals preying less on teleosts (20% IRI) and more on marine mammals (61% IRI) (Kohatsu 2020). In other places (California, USA; South Africa; Argentina; Tasmania, Australia) where Broadnose Sevengill Shark diet has been studied, adults prey mainly on marine mammals (Ebert 2002; Lucifora et al. 2005; Barnett et al. 2010; Hammerschlag et al. 2019). This area overlaps with large rookeries of South American Sea Lion *Otaria flavescens* located in Lobos de Tierra and Lobos de Afuera Islands, representing the main colonies of pinnipeds in northern Peru (Majluf & Trillmich 1981). This area also overlaps with an Important Marine Mammal Area which represents an ideal habitat for several marine mammals, in particular small cetaceans (i.e., Burmeister's Porpoise *Phocoena spinipinnis* and Dusky Dolphin *Lagenorhynchus obscurus posidonia*) (IMMA 2022). Furthermore, the most important landing sites in Peru for Broadnose Sevengill Shark is adjacent to this area (i.e., Lambayeque region) (IMARPE landings statistics).

Juvenile Chilean Eagle Ray prey on teleost fishes (Peruvian Hake and Peruvian Anchoveta), crustaceans (i.e., crabs and stomatopods), gastropod molluscs, and polychaetes (Torres 1978; Castañeda 1994; Segura-Cobeña 2017; Peña-Cutimbo et al. 2025). Thus, this species feeds on both pelagic and demersal prey. Most recent studies sampled 245 individuals and found 94% (in 2015) and 65% (in 2021) of stomachs contained food items. Prey varied according to body size, seasonality, and ENSO conditions in which the Peruvian Anchoveta represented an important prey, especially during warmer conditions (Gonzalez-Pestana et al. 2021b; Peña-Cutimbo et al. 2025).


An analysis of stomach contents from 74 Pacific Guitarfish (mostly adults) between January 2015 and August 2016, indicated that prey included coastal crustaceans (stomatopods and crabs) and teleosts (mainly Peruvian Anchoveta); diet varied according to ontogeny (Gonzalez-Pestana et al. 2021a, 2021b).

SUB-CRITERION D2 – DIVERSITY

Piura-La Libertad sustains a high diversity of Qualifying Species (17 species). This equals the regional diversity threshold (17 species) for the Central and South Pacific American region.

Pelagic Thresher, Common Thresher, Diamond Stingray, and Spinetail Devil Ray are frequently captured by small-scale fisheries operating and landing in this area (Gonzalez-Pestana et al. 2016, 2022; Alfaro-Cordova et al. 2017; Torres 2017; Gonzalez-Pestana 2022). Other species such as the Rasptail Skate is the most abundant incidentally captured ray species (e.g., a total of 556 individuals captured between April and July of 2009) and represents the highest catch-per-unit-effort in the trawl fishery for Peruvian Hake that operates within this area (Céspedes 2014). Large female adults

of Whale Sharks aggregate along the shelf break in this area from December through March (Hearn et al. 2016, 2017; Ryan et al. 2017).



Acknowledgments

Ximena Velez-Zuazo (Smithsonian Institute), Joanna Alfaro-Shigueto (ProDelphinus; Universidad Científica del Sur), Francisco Córdova-Zavaleta (Centro Interdisciplinario de Ciencias Marinas), and Adriana Gonzalez Pestana (IUCN SSC Shark Specialist Group - ISRA Project; ProDelphinus) contributed and consolidated information included in this factsheet. We thank the participants of the 2022 ISRA Region 12 - Central and South American Pacific workshop for their contributions to this process.

This factsheet has undergone review by the ISRA Independent Review Panel prior to its publication.

This project was funded by the Shark Conservation Fund, a philanthropic collaborative pooling expertise and resources to meet the threats facing the world's sharks and rays. The Shark Conservation Fund is a project of Rockefeller Philanthropy Advisors.

Suggested citation

IUCN SSC Shark Specialist Group. 2026. Piura-La Libertad ISRA Factsheet. Dubai: IUCN SSC Shark Specialist Group.

QUALIFYING SPECIES

Scientific Name	Common Name	IUCN Red List Category	Global Depth Range (m)	ISRA Criteria/Sub-criteria Met								
				A	B	C1	C2	C3	C4	C5	D1	D2
SHARKS												
<i>Alopias pelagicus</i>	Pelagic Thresher	EN	0-300	X								X
<i>Alopias vulpinus</i>	Common Thresher	VU	0-150	X								
<i>Carcharhinus brachyurus</i>	Copper Shark	VU	1-145	X		X	X					
<i>Galeorhinus galeus</i>	Tope Shark	CR	0-826	X		X	X					
<i>Mustelus whitneyi</i>	Humpback Smoothhound	CR	16-211	X	X	X	X					
<i>Notorynchus cepedianus</i>	Broadnose Sevengill Shark	VU	0-570	X			X					
<i>Rhincodon typus</i>	Whale Shark	EN	0-1,928	X								
<i>Sphyrna zygaena</i>	Smooth Hammerhead	VU	1-200	X		X	X					
<i>Squatina armata</i>	Chilean Angelshark	CR	0-400	X	X							
<i>Triakis maculata</i>	Spotted Houndshark	CR	10-200	X	X		X					


QUALIFYING SPECIES


Scientific Name	Common Name	IUCN Red List Category	Global Depth Range (m)	ISRA Criteria/Sub-criteria Met								
				A	B	C1	C2	C3	C4	C5	D1	D2
RAYS												
<i>Hypanus dipterurus</i>	Diamond Stingray	VU	0-150	X								X
<i>Mobula mobular</i>	Spinetail Devil Ray	CR	0-1112	X								
<i>Myliobatis chilensis</i>	Chilean Eagle Ray	VU	0-100	X	X	X	X					
<i>Myliobatis peruvianus</i>	Peruvian Eagle Ray	VU	0-50	X	X							
<i>Pseudobatos planiceps</i>	Pacific Guitarfish	VU	1-50	X			X					
<i>Rostroraja velezi</i>	Rasptail Skate	VU	30-300	X								
<i>Sympterygia brevicaudata</i>	Shorttail Fanskate	NT	8-100		X							

SUPPORTING SPECIES

Scientific Name	Common Name	IUCN Red List Category
SHARKS		
<i>Echinorhinus cookei</i>	Prickly Shark	DD
<i>Heterodontus quoyi</i>	Galápagos Bullhead Shark	LC
<i>Megachasma pelagios</i>	Megamouth Shark	LC
RAYS		
<i>Gymnura crebripunctata</i>	Mazatlán Butterfly Ray	NT
<i>Pteroplatytrygon violacea</i>	Pelagic Stingray	LC
<i>Tetronarce tremens</i>	Chilean Torpedo	LC
<i>Urotrygon chilensis</i>	Blotched Round Ray	NT

IUCN Red List categories: *CR*, Critically Endangered; *EN*, Endangered; *VU*, Vulnerable; *NT*, Near Threatened; *LC*, Least Concern; *DD*, Data Deficient.





REFERENCES

- Alfaro-Shigueto J, Mangel JC, Pajuelo M, Dutton PH, Seminoff JA, Godley BJ. 2010.** Where small can have a large impact: structure and characterization of small-scale fisheries in Peru. *Fisheries Research* 106: 8–17. <https://doi.org/10.1016/j.fishres.2010.06.004>
- Alfaro-Cordova E, del Solar A, Alfaro-Shigueto J, Mangel JC, Diaz B, Carrillo O, Sarmiento D. 2017.** Captures of manta and devil rays by small-scale gillnet fisheries in northern Peru. *Fisheries Research* 195: 28–36. <https://doi.org/10.1016/j.fishres.2017.06.012>
- Arellano CE, Swartzman G. 2010.** The Peruvian artisanal fishery: changes in patterns and distribution over time. *Fisheries Research* 101: 133–145.
- Bakun A, Weeks SJ. 2008.** The marine ecosystem off Peru: What are the secrets of its fishery productivity and what might its future hold? *Progress in Oceanography* 79: 290–299. <https://doi.org/10.1016/j.pocean.2008.10.027>
- Barnett A, Abrantes K, Stevens JD, Yick JL, Frusher SD, Semmens JM. 2010.** Predator-prey relationships and foraging ecology of a marine apex predator with a wide temperate distribution. *Marine Ecology Progress Series* 416: 189–200. <https://doi.org/10.3354/meps08778>
- Bertrand A, Segura M, Gutiérrez M, Vasquez L. 2004.** From small-scale habitat loopholes to decadal cycles: a habitat-based hypothesis explaining fluctuation in pelagic fish populations of Peru. *Fish and Fisheries* 5: 296–316.
- Bolaño N. 2009.** Ecología trófica del tiburón martillo *Sphyrna zygaena* (Linnaeus, 1758) en aguas Ecuatorianas. Unpublished Master's Thesis, Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, La Paz.
- Carbajal W, Castañeda J, Castro J, De la Cruz J, Galán J, Ramírez P, Bances S, Salcedo J, Rojas VM. 2007.** Seguimiento e investigación de las pesquerías artesanales en Lambayeque. Informe Anual del Centro de Investigaciones Pesqueras de Santa Rosa del Instituto del Mar del Perú.
- Castañeda J. 1994.** La pesquería artesanal y biología pesquera de especies de importancia económica en la caleta San Jose desde Feb. 1991 a Set. 1992. Unpublished Bachelor Thesis, Universidad Nacional Mayor de San Marcos, Lima.
- Castañeda J. 2001.* Biología y pesquería del tiburón martillo (*Sphyrna zygaena*) en Lambayeque, 1991–2000. *Informe Progresivo IMARPE* 139: 17–31.
- Castillo PR, Madureira L, Marangoni J, Gerlotto F, Guevara-Carrasco R. 2015.** Variability in distribution and aggregation behavior of the Peruvian anchovy (*Engraulis ringens*) analyzed using a fifteen year long series of acoustic surveys (2000–2014). In 2015 IEEE/OES Acoustics in Underwater Geosciences Symposium (RIO Acoustics) (pp. 1–9). IEEE.
- Convention on Biological Diversity (CBD). 2026.** Ecologically or Biologically Significant Marine Areas (EBSAs). Available at: <https://www.cbd.int/ebsa/repository> Accessed January 2026
- Chavez FP, Bertrand A, Guevara-Carrasco R, Soler P, Csirke J. 2008.** The Northern Humboldt Current System: brief history, present status and a view towards the future. *Progress in Oceanography* 79: 95–105.
- Chavez FP, Messie M. 2009.** A comparative analysis of eastern boundary upwelling ecosystems. *Progress in Oceanography* 83: 80–96. <https://doi.org/10.1016/j.pocean.2009.07.032>
- Cliff G, Dudley SFJ. 1992.** Sharks caught in the protective gill nets off Natal, South Africa. 6. The copper shark *Carcharhinus brachyurus* (Günther). *South African Journal of Marine Science* 12: 663–674.
- Céspedes C. 2014.** Captura incidental de fauna marina en la pesquería de *Merluccius gayi peruanus*: Análisis y recomendaciones para su mitigación. Unpublished Master Thesis, Universidad Nacional Agraria la Molina, Lima.
- Córdova-Zavaleta, FA. 2022.** La pesquería artesanal de elasmobranchios en la zona norte-centro del Perú, redes de enmalle como caso de estudio. Unpublished Master Thesis, Instituto Politecnico Nacional, La Paz.
- Csirke J, Argüelles Torres J, Alegre Norza Sior ARP, Ayón Dejo P, Bouchon Corrales M, Castillo**

Mendoza G, Castillo R, Cisneros R, Guevara-Carrasco R, Lau L, et al. 2018. Biología, estructura poblacional y pesquería de pota o calamar gigante (*Dosidicus gigas*) en el Perú. *Boletín Instituto del Mar del Perú* 33: 302-364.

Secretariat of the Convention on Biological Diversity (CBD). 2020. Ecologically or Biologically Significant Marine Areas (EBSAs). Special places in the world's oceans. Volume 5: Eastern Tropical and Temperate Pacific Ocean. Available at: <https://www.cbd.int/marine/ebsa/booklet-05-ettp-en.pdf> Accessed September 2022.

Drew M, Rogers P, Huveneers C. 2017. Slow life-history traits of a neritic predator, the bronze whaler (*Carcharhinus brachyurus*). *Marine and Freshwater Research* 68: 461-472. <https://doi.org/10.1071/MF15399>

Ebert DA. 2002. Ontogenetic changes in the diet of the sevengill shark (*Notorynchus cepedianus*). *Marine and Freshwater Research* 53: 517-523. <https://doi.org/10.1071/MFO1143>

Ebert DA, Dando M, Fowler S. 2021. *Sharks of the World: A complete guide*. Plymouth: Wild Nature Press.

Estupiñan-Montaño C, Cedeño-Figueroa LG, Galván-Magaña F. 2009. Hábitos alimentarios del tiburón martillo *Sphyrna lewini* (Griffith & Smith, 1834) (Chondrichthyes) en el Pacífico ecuatoriano. *Revista de Biología Marina y Oceanografía* 44: 379-386.

FAO. 2022. The state of world fisheries and aquaculture 2022. Towards blue transformation. Rome: FAO. Available at: <https://doi.org/10.4060/cc0461en> Accessed October 2022.

Galván-Magaña F, Polo-Silva C, Hernández-Aguilar SB, Sandoval-Londoño A, Ochoa-Díaz MR, Aguilar-Castro N., Castañeda-Suarez D, Cabrera A, Baigorri-Santacruz A, Torres-Rojas YE, Abitia-Cárdenas LA. 2013. Shark predation on cephalopods in the Mexican and Ecuadorian Pacific Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 95: 52-62.

Garnique-Capuñay D, Jimenez A, Pingo S, Peña-Cutimbo N, Alfaro-Shigueto J, Mangel JC. 2025. New insights into the diet and feeding ecology of the South Pacific angelshark, *Squatina armata* (Philippi, 1887) from Peru. *Journal of Fish Biology*. <https://doi.org/10.1111/jfb.70291>

Graham W, Field J, Potts D. 1992. Persistent 'upwelling shadows' and their influence on zooplankton distributions. *Marine Biology* 114: 561-570.

Gonzalez-Pestana A. 2014. Ecología trófica y áreas de crianza del tiburón martillo, *Sphyrna zygaena* (Linnaeus 1758), juvenil de la zona del norte del Perú. Unpublished Bachelor Thesis, Universidad Científica del Sur, Lima.

Gonzalez-Pestana A, Kouri C, Velez-Zuazo X. 2016. Shark fisheries in the Southeast Pacific: A 61-year analysis from Peru. *F1000Research* 3. <https://doi.org/10.12688/f1000research.4412.2>

Gonzalez-Pestana A, Acuna-Perales N, Coasaca-Céspedes J, Cordova-Zavaleta F, Alfaro-Shigueto J, Mangel JC, Espinoza P. 2017. Trophic ecology of the smooth hammerhead shark (*Sphyrna zygaena*) off the coast of northern Peru. *Fishery Bulletin* 115: 451-459.

Gonzalez-Pestana A. 2018. Habitat suitability of juvenile smooth hammerhead shark (*Sphyrna zygaena*) off northern Peru. Unpublished Master Thesis, James Cook University, Queensland.

Gonzalez-Pestana A, Alfaro-Shigueto, J, Mangel, J. 2019. Aspects of reproductive biology of the humpback smooth-hound shark (*Mustelus whitneyi*) off northern Peru. *Marine and Freshwater Research* 70: 1185-1188. <https://doi.org/10.1071/MF18382>

Gonzalez-Pestana A, Mangel JC, Alfaro-Córdova E, Acuña-Perales N, Córdova-Zavaleta F, Segura-Cobeña E, Benites D, Espinoza M, Coasaca-Céspedes J, Jiménez A, et al. 2021a. Diet, trophic interactions and possible ecological role of commercial sharks and batoids in northern Peruvian waters. *Journal of Fish Biology* 98: 768-783.

Gonzalez-Pestana A, Silva-Garay L, Quiñones J, Mayaute L, Manrique M, Segura-Cobeña E, Espinoza P, Moscoso V, Velez-Zuazo X, Alfaro-Shigueto J, Mangel JC. 2021b. Geographic and ontogenetic variation in the diet of two commonly exploited batoids (Chilean eagle ray and Pacific guitarfish) off Peru: evidence of trophic plasticity. *Environmental Biology of Fishes* 104: 1525-1540. <https://doi.org/10.1007/s10641-021-01157-w>

- Gonzalez-Pestana A, Velez-Zuazo X, Alfaro-Shigueto J, Mangel JC. 2022.** Batoid fishery in Peru (1950-2015): Magnitude, management and data needs. *Revista de Biología Marina y Oceanografía*. 57: 217-233. <https://doi.org/10.22370/rbmo.2022.57.Especial.3729>
- Gonzalez-Pestana A. 2022.** Catch composition of mobulid rays (*Mobula* spp.) in northern Peru reveals a potential nursery area for *M. mobular*. *Environmental Biology of Fishes* 105: 963-969. <https://doi.org/10.1007/s10641-022-01301-0>
- Gonzalez-Pestana A, Alfaro-Shigueto J, Mangel JC. 2022.** A review of high trophic predator-prey relationships in the pelagic Northern Humboldt system, with a focus on anchovetas. *Fisheries Research* 253: 106386. <https://doi.org/10.1016/j.fishres.2022.106386>
- Hammerschlag N, Williams L, Fallows M, Fallows C. 2019.** Disappearance of white sharks leads to the novel emergence of an allopatric apex predator, the sevengill shark. *Scientific Reports* 9: 1908. <https://doi.org/10.1038/s41598-018-37576-6>
- Hearn A, J Green, MH Roman, D Acuña-Marrero, E Espinoza, Klimley AP. 2016.** Adult female whale sharks make long distance movements past Darwin Island (Galapagos, Ecuador) in the Eastern Tropical Pacific. *Marine Biology* 163: 213-224. <https://doi.org/10.1007/s00227-016-2991-y>
- IUCN. 2025.** The IUCN Red List of Threatened Species. Version 2025-1, Available at: <https://www.iucnredlist.org/> Accessed November 2025.
- Important Marine Mammals Areas (IMMA). 2022.** Continental Shelf of the Northern Humboldt Current IMMA. Available at: <https://www.marinemammalhabitat.org/portfolio-item/continental-shelf-of-the-northern-humboldt-current-imma/> Accessed December 2022.
- Jacox MG, Edwards CA. 2011.** Effects of stratification and shelf slope on nutrient supply in coastal upwelling regions. *Journal of Geophysical Research: Oceans* 116: C03019. <https://doi.org/10.1029/2010JC006547>
- Jian W, Jingwen G, Ting L, Songling Z, Yuanyuan T, Xinjun C, Wei Y. 2020.** Spatio-temporal variations in the habitat of jumbo squid *Dosidicus gigas* in the Southeast Pacific Ocean off Peru under anomalous climate conditions. *Haiyang Xuebao* 42: 92-99.
- Key Biodiversity Areas (KBA). 2026.** Key Biodiversity Areas factsheet: Sistema de Islas, Islotes y Puntas Guaneras. Available at: <https://www.keybiodiversityareas.org/site/factsheet/200718> Accessed February 2026.
- Kohatsu SN. 2020.** Ecología trófica del tiburón de siete agallas *Notorynchus cepedianus* en el norte del Perú. Unpublished Bachelor Thesis, Universidad Científica del Sur, Lima.
- Llanos J, Galán J, Castañeda J, Castro J, De la Cruz J, Ramírez P, Bances S, Torres D. 2009.** Investigaciones de IMARPE Sede Lambayeque Durante 2009. Reporte de Investigación del Instituto del Mar del Perú (IMARPE).
- Lucifora LO, Menni RC, Escalante AH. 2005.** Reproduction, abundance and feeding habits of the broadnose sevengill shark *Notorynchus cepedianus* in north Patagonia, Argentina. *Marine Ecology Progress Series* 289: 237-244. <https://doi.org/10.3354/meps289237>
- Lucifora LO, García VB, Menni RC, Escalante AH, Hozbor NM. 2009.** Effects of body size, age and maturity stage on diet in a large shark: ecological and applied implications. *Ecological Research* 24: 109-118. <https://doi.org/10.1007/s11284-008-0487-z>
- Majluf P, Trillmich F. 1981.** Distribution and abundance of sea lions (*Otaria byronia*) and fur seal (*Arctocephalus australis*) in Peru. *Zeitschrift Fur Säugetierkunde* 46: 384-393.
- Montecino V, Lange CB. 2009.** The Humboldt Current System: Ecosystem components and processes, fisheries, and sediment studies. *Progress in Oceanography* 83: 65-79. <https://doi.org/10.1016/j.pocean.2009.07.041>
- Pennington JT, Mahoney KL, Kuwahara VS, Kolber DD, Calienes R, Chavez FP. 2006.** Primary production in the eastern tropical Pacific: a review. *Progress in Oceanography* 69: 285-317. <https://doi.org/10.1016/j.pocean.2006.03.012>

- Peña-Cutimbo N, Cordova-Zavaleta F, Velarde-Vilchez M, Jimenez A, Pingo S, Ramirez P, Castaneda J, De la Cruz J, Mange JC, Alfaro-Shigueto J. 2025. Comparative feeding analysis between the Chilean eagle ray *Myliobatis chilensis* (Philippi, 1982) and the Peruvian eagle ray *Myliobatis peruvianus* (Garman, 1913) from northern Peru. *Regional Studies in Marine Science* 81: 103915. <https://doi.org/10.1016/j.rsma.2024.103915>
- Peres MB, Vooren CM. 1991. Sexual development, reproductive cycle, and fecundity of the school shark *Galeorhinus galeus* off southern Brazil. *Fishery Bulletin* 89: 655-667.
- Ryan JP, Green JR, Espinoza E, Hearn AP. 2017. Association of whale sharks (*Rhincodon typus*) with thermo-biological frontal systems of the eastern tropical Pacific. *PLoS One* 12: e0182599. <https://doi.org/10.1371/journal.pone.0182599>
- Samame M, Castillo J, Espino M. 1989. El tollo un recurso demersal, algunos aspectos de la biología y pesquería de *Mustelus whitneyi*. *Revista del Pacifico Sur*. Special Volume 1989
- Segura-Cobeña E. 2017. Composición de la dieta de la Raya Águila Chilena *Myliobatis chilensis* (Philippi, 1892) en la caleta de San José, Lambayeque. Unpublished Bachelor Thesis, Universidad Científica del Sur, Lima.
- Sencio-Sánchez DC. 2025. Aspectos biológico-pesqueros de *Squatina armata* (angelote), caleta San José Lambayeque, julio 2022 a abril 2023. Unpublished Bachelor Thesis. Universidad Nacional Pedro Ruiz Gallo, Lambayeque.
- Smale MJ. 1991. Occurrence and feeding of three shark species, *Carcharhinus brachyurus*, *C. obscurus* and *Sphyrna zygaena*, on the Eastern Cape coast of South Africa. *South African Journal of Marine Science* 11: 31-42.
- Torres Carrasco AA. 2018. Variación espacio-temporal en las capturas provenientes de la pesquería artesanal de tiburón con red de enmalle de superficie, durante octubre 2016 a marzo 2018. Unpublished Bachelor Thesis, Universidad Nacional de Piura, Piura.
- UNEP-WCMC & IUCN. 2026. Protected Planet: The World Database on Protected Areas (WDPA) and World Database on Other Effective Area-based Conservation Measures (WD-OECM) [Online], February 2026, Cambridge, UK: UNEP-WCMC and IUCN. Available at: www.protectedplanet.net. Accessed February 2026.
- Vargas N, Mendo J. 2010. Relación entre la distribución espacial de la merluza peruana (*Merluccius gayi peruanus* Ginsburg) y la extensión sur de la corriente de Cromwell. *Informe IMARPE* 37(3-4).
- Zuta S, Guillen O. 1970. Oceanografía de las aguas costeras del Perú. *Boletín Instituto del Mar del Perú* 2: 157-324.